

Rhyme® Fungicide

RHYME™

FUNGICIDE

Highly Systemic for Maximum Plant Protection

As a rotational partner against powdery mildew during rapid vine growth, you would be hard pressed to make a better Group 3 fungicide selection than Rhyme™ fungicide. Rhyme fungicide protects both crop quality and yields.

Rhyme fungicide is one of the most systemic fungicides in its class, moving throughout the plant faster than other triazoles and offering full protection where other leading fungicides fail. The quick penetration and rapid movement to sites of established infection provides fast curative activity on powdery mildew. As the most residual of all triazoles, Rhyme fungicide provides longer lasting protection to prevent further infections.

Rhyme fungicide is rainfast within two hours and can be tank mixed with most fungicides, insecticides and surfactants for increased crop protection. Rhyme fungicide is an excellent choice as a foundational fungicide. (Select link below for testimonial.)

<https://www.youtube.com/embed/GbtFypkddME>



Application Guidelines

Maximum rate per acre: 5 fl. Oz.

Re-treatment interval: 14 days

May be applied via ground, air, or chemigation including drip irrigation and micro-sprinklers. (See 2ee label for pests and instructions)

Preharvest Interval (PHI): 14 days

Maximum applications per crop year: 6

Restricted Entry Interval (REI): Re-entry interval for girdling or turning of grapes is 5 days. The REI for all other activities is 12 hrs

Diseases	Application Instructions
Powdery Mildew (<i>Uncinula necator</i>)	Apply Rhyme fungicide beginning at 6- 10 inch shoots. Repeat as necessary if conditions are favorable for disease development.
Black Rot (<i>Guignardia bidwellii</i>)	Apply Rhyme fungicide preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.

Drip Irrigation and Micro-sprinkler Chemigation Use Guidelines

For control of foliar powdery mildew use Rhyme™ fungicide in two consecutive drip or micro-sprinkler applications as part of a standard foliar spray program of registered products.

- Apply preventatively or when conditions are favorable for disease development.
- Apply the first application beginning at 6" - 10" shoot growth
- Apply the second application 14 days following the first application.

Apply in sufficient irrigation water to wet the rooting profile of the grapes or to 12 inches of soil. Delay subsequent irrigation for 24 hours following a drip application or micro-sprinkler application for best results.


Restrictions:

- Do not apply Rhyme fungicide through drip irrigation that is deeper than 12" below the soil surface.
- See label for a complete list of restrictions.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. A person knowledgeable of the chemigation system and responsible for its operation will shut down the system and make necessary adjustments should the need arise. If you have questions about calibration, you should contact state extension service specialist, equipment manufacturer, or other experts.

Foliar Applied Rhyme Fungicide

Percent Powdery Mildew Infected Clusters



Always read and follow label directions. FMC and Rhyme are trademarks and Investing in farming's future is a service mark of FMC Corporation or an affiliate. Mettle is a trademark of Gowan. Rally is a trademarks of Dow AgroSciences LLC. Inspire is a trademark of Syngenta Crop Protectopn, LLC. ©FMC 2017. All Rights Reserved.