


FirstShot® SG Burndown Herbicide With TotalSol® Soluble Granules



Preplant burndown partner

FirstShot® SG herbicide delivers:

- A short residual, non-volatile glyphosate tank-mix partner for enhanced control of curly dock, filaree, henbit, sunflowers, shepherd's purse, wild mustard and many other labeled weeds before planting cotton, corn, soybeans, sorghum, rice, wheat or other crops.
- Improved control of problem weeds.
- Clean seedbed at planting.
- Better stand establishment of the desired crop due to a cleaner seedbed.
- Rotational freedom with short preplant intervals.



OKLAHOMA AND TEXAS

RATE RECOMMENDATION

For control of curly dock, filaree, henbit, marestail, sunflowers, shepherd's purse and wild mustard

FirstShot SG herbicide 0.6-0.8 oz./A
(depending on weed size)
+ Abundit® Extra herbicide (glyphosate)
or
+ 2,4-D LVE

Application information

- Always include a spray adjuvant with FirstShot SG herbicide.
- When applying 0.5 to 0.6 oz./A, FirstShot SG herbicide must be used in combination with another registered burndown herbicide.
- FirstShot SG herbicide should always be the first product added to the clean water in the tank.

Time interval before planting* (days after treatment with FirstShot® SG herbicide)

FirstShot SG herbicide may be applied prior to planting the following crops:

CROP	DAYS
Barley, rice, triticale and wheat including durum	0
Soybeans	7
Cotton, field corn and grain sorghum	14
Any other crop	45

* Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.

Where FirstShot SG herbicide is used on light textured soils, such as sands, loamy sands and sandy loams, extend time to planting by seven additional days.

Where FirstShot SG herbicide is used on high pH soils (>7.9), extend time to planting by seven additional days.

FirstShot SG burndown herbicide

Weeds Controlled

Buckwheat, common	Groundsel, common	Radish, common
Bushy wallflower/Treacle mustard	Henbit*	Radish, wild*
Buttercup, bur	Knawel, annual	Ragweed, common*
Buttercup, smallflower	Knotweed, prostrate	Redmaids
Catchfly, nightflowering	Lady's thumb	Scentless chamomile/mayweed
Chamomile (corn, false, mayweed, wild)	Lambsquarters (common, narrowleaf, slimleaf)	Shepherd's purse
Chickweed, field	Lettuce, prickly‡	Smartweed (green, Pennsylvania)
Chickweed (common, stinking)	London rocket	Sowthistle, annual
Cocklebur, common*	Marshelder	Spurry, corn
Cowcockle	Miner's lettuce	Stinking mayweed/dogfennel
Cress (mouse-ear)	Mustard (black, blue/purple, Tumble/ Jim Hill mustard, wild)	Sunflower, common
Dock (broadleaf, curly)	Pennycress, field	Sunflower, volunteer
False flax, small seed	Pepperweed, clasping	Swinecress
Fiddleneck (coast, tarweed)	Pigweed (prostrate, redroot)	Tansy mustard
Filaree (redstem, Texas)	Pineapple weed	Volunteer lentils
Flixweed		Volunteer peas
Geranium, Carolina		Wild garlic*

Weeds Partially Controlled**

Bedstraw, catchweed	Mallow (common, little)	Nightshade (cutleaf, hairy)
Evening primrose, cutleaf*	Marestail*	Vetch* (common, hairy)

* See SPECIFIC WEED PROBLEMS for more information.

** Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest directed rate of FirstShot SG herbicide per acre and include a tank-mix partner such as 2,4-D, a dicamba containing product, or a glyphosate containing product.

‡ Naturally occurring resistant biotypes of prickly lettuce, marestail and mayweed spp. are known to occur. See the Resistance section of this label for additional details.

For more information, contact your local FMC retailer or representative to learn more about FirstShot SC herbicide from FMC and visit us at FMCCrop.com.

Always read and follow label directions.

FMC, TotalSol, FirstShot and Abundit are trademarks of FMC Corporation or an affiliate. ©2017 FMC Corporation. All rights reserved. 17-FMC-0978 10/17

