

HatchTrakSM



*Controlling Aphids,
Fleahoppers & Plantbugs*

Make Carbine[®] 50WG insecticide Your Choice to Control Aphids, Fleahoppers and Plantbugs

With the Southeast cotton crop starting off under very wet conditions and as temperatures move higher growers should expect insect pest pressure to increase. In the Southeast, cotton growers have already started fighting pests like aphids, fleahoppers and plantbugs. Carbine[®] 50WG insecticide will control these pests and, at the same time have a low impact on beneficial insects. Carbine 50WG insecticide will not flare aphids.

Coverage is critical to achieve control with Carbine 50WG insecticide. At the time of application, if weather conditions turn hot and dry, it is not recommended to tank mix insecticides with herbicides, especially if you are using air induction tips and low pressure to prevent herbicide drift. In addition to using proper spray tips, organosilicone type adjuvants can be used under these conditions to enhance canopy penetration and coverage of Carbine 50WG insecticide.

Inside This Issue

**CARBINE 50WG
INSECTICIDE**

Controls

- ✓ **Aphids**
- ✓ **Fleahoppers**
- ✓ **Plantbugs**

**Contact your local FMC
representative for more
information:**

**Del Brandenburg
Retail Market Manager
229-392-7662**

**Mike Cunnane
Retail Market Manager
901-201-1657**

**Cindy Kurtz
Retail Market Manager
478-955-2344**

**James Marsh
Retail Market Manager
252-702-2041**


**Craig Heim
Technical Services
Manager
856-261-5659**

FMC



CARBINE[®] 50WG INSECTICIDE APPLICATION RECOMMENDATIONS

- Aphids 1.7-2.0 oz./A. Label rates 1.4-2.8 oz./A.
- Fleahoppers and plant bugs: 1.7-2.8 oz./A.
- Allow seven days between applications.
- Thorough spray coverage of plant foliage is essential for optimum control. Using higher application spray volumes will improve coverage. Recommended by ground 10 GPA and 5 GPA by air (minimum 5 GPA by ground and 3 GPA by air).
- Do not use air induction spray tips, such as those used to reduce herbicide drift.
- 30-day PHI; 12-hour REI.
- Carbine 50WG may be used ahead of acephate and will not flare aphids.



888-59-FMC-AG FMCCROP.COM

Always read and follow all label directions, restrictions and precautions for use. Some products may not be registered for sale or use in all states. FMC is a trademark and HatchTrak is a service mark of FMC Corporation or its affiliates. Carbine is a trademark of Ishihara Sangyo Kaisha, Ltd. Assail is a trademark of Nippon Soda Co., Ltd. Admire is a trademark of Bayer. Centric is a trademark of a Syngenta Group Company. Belay is a trademark of Sumitomo Chemical Co. LTD. Transform is a trademark of Dow Agrosiences LLC. Bidrin is a trademark of AMVAC Chemical Corporation. ©2018 FMC Corporation. All rights reserved. 07/18