



# Peanut Weed Control

**Pre-emergence Power of Spartan with  
Added Pre-plant Burndown!**



Spartan® Charge is an effective pre-emergence broadleaf herbicide labeled for use in peanuts. Spartan Charge provides added flexibility for burndown on pre-formed beds that have gone weedy and need to be freshened up prior to planting. Spartan Charge gives consistent, long-lived control of many broadleaf weeds, including morningglories, pigweed, palmer amaranth and nightshade as well as nutsedge suppression. Spartan Charge is an excellent tank mix partner with other grass and broadleaf herbicides labeled for peanuts.

	Features	Benefits
	Unique Modes of Action	Controls shifting weed populations & manages resistance
	Flexible Application	Residual pre-emergence activity with added burndown performance in one shot
	Long Residual Properties and Broad Spectrum	Saves on labor by removing pigweeds and morningglories
	Tank Mixes Easily	Allows growers to choose between various grass herbicides
Controls Tough Broadleaves	Pigweeds, palmer amaranth, morningglories, Pennsylvania smartweed, and nightshade	

Spartan Charge Rate (oz/A)				Rotational Crop Guideline	
Soil Texture					
% OM	Coarse	Medium	Fine		
<1.5%	3.0 to 3.75	4.0	5.0	Soybean, Sunflower, Dry Peas	Anytime
1.5-3%	4.0	5.0	6.0	Tobacco, Turf, Sugarcane	Anytime
>3%	5.0	6.0	7.0	Wheat, Barley, Rye, Triticale, Field Corn, Potato	4 months
				Sorghum, Rice	10 months
				Corn, : Seed., Pop, Sweet	12 months
				Alfalfa, Oats, Sweet Potato	12 months
				Cotton	12 months

Application Precautions	Restrictions
<p><b>Use proper rate based upon soil texture and organic matter</b></p> <ul style="list-style-type: none"> <li>Coarse soils – sand, loamy sand, sandy loam</li> <li>Medium soils – silt, loam, sandy clay loam, sandy clay, silt loam</li> <li>Fine soils – clay, clay loam, silty clay, silty clay loam</li> </ul> <p><b>Use lower rates where the soil pH is greater than 7.0</b></p>	<ul style="list-style-type: none"> <li>Do not apply to soils classified as sands with less than 1% OM (Organic Matter)</li> <li>Do not apply after crop emergence, at cracking or if seedling is close to the soil surface as undesirable crop response may occur</li> <li>Do not perform tillage practices that concentrates Spartan Charge into the bed or crop injury may occur</li> <li>Do not irrigate when peanuts are cracking or with irrigation water having a pH higher than 9.</li> </ul>