

# Aim® EC Herbicide

## Powerful defoliant and harvest aid

### COTTON

#### PRODUCT INFORMATION:

Aim® EC herbicide from FMC is your secret weapon come harvest. It makes cotton defoliation fast, economical and easy. Aim EC herbicide is unsurpassed for cotton defoliation, juvenile regrowth removal and weed desiccation in spindle and stripper cotton.

Aim EC herbicide also delivers fast-acting, non-residual control of the toughest broadleaf weeds, including kochia, waterhemp, lambsquarters, Russian thistle and more. Flexible and easily applied alone or with a tank-mix partner for harvest aid and defoliation.

#### QUICK FACTS:

- Labeled for cotton, row crops and some specialty crops
- Can be tank mixed with other cotton harvest aid products
- Low use rate for cotton  
Up to 1.6 fl. oz./A
- Controls and suppresses broadleaf weeds
- Flexibility to fit into many different defoliation programs
- Effective in desiccating weeds present in cotton crop



## Cotton Harvest Aid/Defoliant

BRAND/CHEMISTRY	REMOVE MATURE FOLIAGE	REMOVE IMMATURE FOLIAGE	REGROWTH SUPPRESSION	JUVENILE REGROWTH REMOVAL	OPEN BOLLS	DESICCATE VINES AND WEEDS
Aim® EC herbicide + adjuvant	G	E	F	E	F	G
Aim EC herbicide + thidiazuron	E	E	E	E	F	G
Aim EC herbicide + ethephon	E	E	F	E	E	G
Aim EC herbicide + thidiazuron + ethephon	E	E	E	E	E	G
Aim EC herbicide + paraquat	E	E	F	E	F	E

= Excellent
  = Good
  = Fair

### Target Weeds:

Cocklebur  
 Hemp sesbania  
 Kochia  
 Lambsquarters  
 Morningglory  
 Nightshade  
 Palmer amaranth  
 Pigweed  
 Russian thistle  
 Velvetleaf  
 Waterhemp  
 Wild buckwheat

### Application Timing:

**Harvest Aid Application:** Apply Aim® EC herbicide alone or as a tank mixture with other cotton harvest aids to defoliate and desiccate cotton and troublesome weeds that may be present at harvest. Use a good quality spray adjuvant at the recommended rates.

### Application Guidelines:

**Pre-Harvest Interval:** Seven days

**Product Use Rate:** 0.25 – 1.6. oz./A

Apply when 60-70 percent of bolls are open or according to the state agricultural extension service guidelines in the use area.

### Restrictions:

Do not apply when conditions favor drift or when wind is above 10 mph. Do not apply more than 7.9 fl. oz./A total for preplant, in-season weed control and harvest aid.

**Please see product label for a complete list of restrictions.**

Do not apply within seven days of harvest.

Re-entry interval (REI): 12 hours

### Label:

**Active Ingredient:** Carfentrazone-ethyl (Group 14), Group E herbicide

**Mode of Action:** The disruption of membranes by inhibiting the action of protoporphyrinogen oxidase, causing cell death.

**To learn more about Aim EC herbicide, contact your FMC Star Retailer or visit [FMCCrop.com](http://FMCCrop.com).**

Always read and follow all label directions, restrictions and precautions for use. Aim EC herbicide is not registered for use in California. FMC and Aim are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation. All rights reserved. 18-FMC-0206 02/18

