


BROAD-SPECTRUM, ECONOMICAL, EARLY AND LATE SEASON INSECT CONTROL

Stink Bug Control



On-Farm Grower Trials

	Treatment	Yield Bu/A
Standard Program	Delayed burndown <i>fb</i> glyphosate	63.6
Intensively Managed	Authority® First DF <i>fb</i> post glyphosate + Mustang® Maxx + Headline®	68.7



Untreated vs. Mustang Maxx - 26 DAT

Application Recommendations

1.28-4.0 oz/acre Cutworm, thistle caterpillar, silverspotted skipper.

2.8-4.0 oz/acre Alfalfa caterpillar, southern armyworm, true armyworm, yellowstriped armyworm, bean leaf beetle, blister beetle, Colorado potato beetle, cucumber beetle, European corn borer, corn earworm, corn rootworm beetle (adult), flea beetle, green cloverworm, hornworms, Japanese beetle, leafhopper, leafminers (adults), Mexican bean beetle, potato leafhopper, seedcorn maggot (adult), soybean aphid, tobacco budworm, three-cornered alfalfa hopper, webworm, woolly bear caterpillar.

3.2-4.0 oz/acre Beet armyworm, fall armyworm, grasshopper, looper, stink bugs, thrips, whitefly.

Special Information **12 hour REI, 21 day PHI.** Apply using sufficient carrier to obtain thorough coverage of foliage to achieve control, minimum of 10 gals/acre by ground or 2 gals/acre by air (include COC or equivalent). Can be tank mixed with commonly used herbicides, fungicides, and foliar fertilizers. Do not make applications less than 7 days apart. Do not graze or harvest treated soybean forage, straw, or hay for livestock feed. Do not apply more than 24 oz/acre per season.

