

## CONTROLS UNIQUE SPECTRUM

- Proven oxadiazine class of chemistry in IRAC Group 22
- Improved formulation for better mixability and enhanced residual
- Expanded label allows higher application rates in certain crops, plus new crop additions including beans, low-growing berries and small fruit vine climbing subgroups
- Non restricted use pesticide
- Effective control of the lepidopteran spectrum, plus several species of weevils, beetles and sucking pests
- Resistant to washoff and stable under UV light and high heat
- Conforms to IPM program best practices
- No mite flares

## PESTS CONTROLLED INCLUDE

- Corn earworm
- European corn borer
- Armyworms
- Cabbage looper
- Cabbage webworm
- Hornworms
- Loopers
- Japanese beetle
- Leafroller
- Leafhoppers
- Spotted cutworm

## SPECIAL INFORMATION

- Apply follow-up treatments as needed, every 3-5 days on most labeled crops
- Please see label information for crop-specific seasonal application limits
- May be applied by ground, aerial or overhead sprinkler chemigation
- For aerial application, use a minimum of 5 GPA and maximum of 200 GPA water unless otherwise specified in label directions.
- Do not use adjuvants on bushberries or garden beets
- PHI: 3 – 7 days\*
- REI: 12 hours
- Use rate: 2.5 – 6 fl. oz./A
- Signal word: Caution

\*Except for stone fruit, pome fruit and labeled low-growing berry subgroup

## NEW FORMULATION FOR BETTER MIXABILITY



Left: Avaunt® insecticide after water is poured over product.  
Right: Avaunt® eVo insect control after water is poured over product.

Always read and follow label directions and precautions for use. Some products may not be registered for sale or use in all states. As of November 1, 2017, the USEPA registrations for DuPont™ Avaunt® insecticide and Avaunt® eVo, Altacor®, Coragen®, Exirel®, Prevathon® and Verimark® insect controls were sold by E.I. du Pont de Nemours and Company to FMC Corporation. FMC, Avaunt, Altacor, Coragen, Exirel, Prevathon and Verimark are trademarks of FMC Corporation or an affiliate. ©2018 FMC Corporation. All rights reserved. 18-FMC-0590 04/18

LABELED CROPS	USE IN ROTATION WITH:
<b>Beans, dried and succulent (except soy)</b>	Coragen® insect control Prevathon® insect control Exirel® insect control
<b>Brassica (cole)</b>	Coragen insect control Verimark® insect control Exirel insect control
<b>Bushberries</b>	Altacor® insect control Exirel insect control
<b>Sweet corn (for application through tassel push only)</b>	Coragen insect control
<b>Low-growing berry subgroup (except strawberries)</b>	Altacor insect control
<b>Cucurbit vegetables</b>	Coragen insect control Verimark insect control Exirel insect control
<b>Fruiting vegetables and okra</b>	Coragen insect control Verimark insect control Exirel insect control
<b>Grapes</b>	Altacor insect control
<b>Small fruit vining subgroup (except fuzzy kiwi)</b>	Altacor insect control
<b>Leafy green vegetables e.g., spinach</b>	Coragen insect control Verimark insect control Exirel insect control
<b>Leafy petioles</b>	Coragen insect control Verimark insect control Exirel insect control
<b>Mint (peppermint and spearmint)</b>	Coragen insect control
<b>Pome fruit</b>	Altacor insect control Exirel insect control*
<b>Stone fruit</b>	Altacor insect control Exirel insect control
<b>Tuberous and corm vegetables e.g., garden beets</b>	Exirel insect control Coragen insect control

\*Use of Exirel insect control and captan may result in adverse crop response. Avoid using Exirel insect control within 14 days of a captan fungicide application.